

## Forests, species on four continents threatened by palm oil expansion

Local conservation strategies, tailored to each region's differing threats, will be needed to protect at-risk species

**Date:** July 27, 2016

**Source:** Duke University

**Summary:** As palm oil production expands from Southeast Asia into the Americas and Africa, vulnerable tropical forests and species on four continents face increased risk of loss, a new study finds. The largest areas of vulnerable forest are in Africa and South America. But because forests in all 20 countries studied contain high concentrations of different mammal and bird species at risk of extinction, conservation efforts need to incorporate localized solutions tailored to each region.

### FULL STORY

---

As palm oil production expands from Southeast Asia into tropical regions of the Americas and Africa, vulnerable forests and species on four continents face increased risk of loss, a new Duke University-led study finds.

The largest areas of vulnerable forest are in Africa and South America, where more than 30 percent of forests within land suitable for oil palm plantations remain unprotected, the study shows.

Rates of recent deforestation have been highest in Southeast Asia and South America, particularly Indonesia, Ecuador and Peru, where more than half of all oil palms are grown on land cleared since 1989.

"Almost all oil palm is grown in places that once were tropical forests. Clearing these forests threatens biodiversity and increases greenhouse gas emissions," said Varsha Vijay, a doctoral student at Duke's Nicholas School of the Environment who led the study.

"By identifying where the greatest extent of palm oil-driven deforestation has recently occurred and modelling where future expansion is most likely, we can guide efforts to reduce these adverse impacts," Vijay said.

Palm oil is now the world's most widely traded vegetable oil, according to the Food and Agriculture Organization of the United Nations. The oil, which is harvested from oil palms, and its derivatives are common ingredients in many processed foods and personal care products. As global demand grows, large swaths of tropical forest are being converted into oil palm plantations in 43 countries.

Using 25 years of high-resolution Google Earth and Landsat satellite imagery, Vijay and her team tracked the extent of this deforestation in four regions: Southeast Asia, Africa, South America and Mesoamerica, which includes Mexico, Central America and the Caribbean.

They modeled where future deforestation is most likely to occur, based on the pace of recent expansion in each region and the extent of suitable forest left unprotected there.

The team published its findings July 27 in the peer-reviewed, open-access journal *PLOS ONE*.

Because forests in all four regions contain high concentrations of different mammal and bird species at risk of extinction or extirpation, conservation efforts will need to incorporate strategies tailored to each region, said Stuart Pimm, Doris Duke Professor of Conservation at Duke.

"While the Amazon and Indonesia have many species of globally threatened mammals and birds, other areas such as the Congo Basin and the coastal forests of Colombia are home to species with small ranges that make them especially vulnerable to habitat loss despite not being classified as threatened or endangered," he said. "This needs to be considered in conservation planning."

"Many past studies have focused solely on Indonesia and Malaysia, which produce more than 80 percent of the world's palm oil. By evaluating deforestation caused by palm oil production in 20 countries across four regions, our study demonstrates that the biodiversity impacts of this expansion are very different from country to country and region to region," said Clinton Jenkins of the Institute for Ecological Research in Brazil.

"The palm oil industry has a legacy of deforestation, and today consumer pressure is pushing companies toward deforestation-free sources of palm oil," noted Sharon Smith of the Union of Concerned Scientists, who co-authored the study with Vijay, Pimm and Jenkins.

"This research helps us understand where to focus on using government regulation and voluntary market interventions to shape oil palm plantation expansion in ways that protect biodiversity-rich ecosystems and prevent deforestation," Smith said.

---

#### Story Source:

The above post is reprinted from materials provided by **Duke University**. *Note: Materials may be edited for content and length.*

---

#### Journal Reference:

1. Varsha Vijay, Stuart L. Pimm, Clinton N. Jenkins, Sharon J. Smith. **The Impacts of Oil Palm on Recent Deforestation and Biodiversity Loss**. *PLOS ONE*, 2016; 11 (7): e0159668 DOI: 10.1371/journal.pone.0159668

---

#### Cite This Page:

MLA	APA	Chicago
-----	-----	---------

---

Duke University. "Forests, species on four continents threatened by palm oil expansion: Local conservation strategies, tailored to each region's differing threats, will be needed to protect at-risk species." ScienceDaily. ScienceDaily, 27 July 2016. <[www.sciencedaily.com/releases/2016/07/160727172011.htm](http://www.sciencedaily.com/releases/2016/07/160727172011.htm)>.

---

#### Recommended Articles

Oil Palm Plantations Are No Substitute For Tropical Rainforests, New Study Shows  
Queen Mary, University of London, ScienceDaily, 2008

Biofuel Plantations On Tropical Forestlands Are Bad For The Climate And Biodiversity, Study

Building on Two Decades of Ecosystem Management and Biodiversity Conservation under the Northwest Forest Plan, USA  
DellaSala, Dominick A.; Baker, Rowan ; Heiken, Doug ; Frissell, Chris A.; Karr, James R.; Nelson, S. Kim; Noon, Barry R.; Olson, David ; Strittholt, James et al., Forests, 2015

## Finds

World Wildlife Fund, ScienceDaily, 2008

Threatened frogs palmed off as forests disappear

Zoological Society of London, ScienceDaily, 2013

Carbon Payments Help Protect Threatened Tropical Mammals

University of Queensland, ScienceDaily, 2009

Biofuels And Biodiversity Don't Mix, Ecologists Warn

Wiley-Blackwell, ScienceDaily, 2008

Scientists Track Seasonal Flu's Global Journey

Catharine Paddock PhD, Medical News Today

Assessment and control of vitamin A deficiency disorders.

Usha Ramakrishnan et al., J Nutr, 2002

The type of oil used for cooking is associated with the risk of nonfatal acute myocardial infarction in costa rica.

Edmond K Kabagambe et al., J Nutr, 2005

Comparison of the effects of supplemental red palm oil and sunflower oil on maternal vitamin A status

Georg Lietz et al., Am J Clin Nutr, 2001

---

Powered by